

## VOLATILITY CONVERSION CALCULATORS

PATRICK S. HAGAN  
 BLOOMBERG  
 499 PARK AVENUE  
 NEW YORK, NY 10022  
 PHAGAN1@BLOOMBERG.NET

**Abstract.** We provide the formulas needed to translate from absolute (normal) volatilities to Black (log normal) volatilities. For good measure, we then give the formulas to convert CEV volatilities into Black and absolute volatilities

**Key words.** equivalent vol, deterministic vol, smile

### 1. Conversion between log normal and normal vol. Black's model is

$$(1.1a) \quad dF = \sigma_B F dW, \quad F(0) = f$$

where  $f$  is today's forward swap/caplet rate and where  $\sigma_B$  is the implied Black (log normal) volatility. The normal model is

$$(1.1b) \quad dF = \sigma_N dW, \quad F(0) = f$$

where  $\sigma_N$  is the "normal" or "absolute" or "bps per year" volatility.

**1.1. Calculating normal vol from log normal vol.** For a swaption with strike (fixed rate)  $K$ , the normal volatility  $\sigma_N$  (which gives the same price of the option) as the log normal volatility  $\sigma_B$  is:

$$(1.2a) \quad \sigma_N = \sigma_B \frac{f - K}{\log f/K} \cdot \frac{1}{1 + \frac{1}{24} \left(1 - \frac{1}{120} \log^2 f/K\right) \sigma_B^2 \tau + \frac{1}{5760} \sigma_B^4 \tau^2}$$

where

$$(1.2b) \quad f = \text{current forward swap or caplet rate,}$$

$$(1.2c) \quad K = \text{option's strike (fixed rate),}$$

$$(1.2d) \quad \tau = \text{time to exercise (notification) date in years,}$$

We probably should dispense with the

$$(1.3) \quad \frac{1}{120} \log^2 f/K \quad \text{and} \quad \frac{1}{5760} \sigma_B^4 \tau^2.$$

They are too small to add measurably to the accuracy.

When  $f \rightarrow K$ , the above formula goes to a "0 over 0." To avoid this complication, we should use the alternative formula

$$(1.4a) \quad \sigma_N = \sigma_B \sqrt{fK} \frac{1 + \frac{1}{24} \log^2 f/K}{1 + \frac{1}{24} \sigma_B^2 \tau + \frac{1}{5760} \sigma_B^4 \tau^2}$$

when

$$(1.4b) \quad \left| \frac{f - K}{K} \right| < 0.001.$$

**1.2. Calculating normal vol from log normal vol.** Now suppose we are given an absolute (normal) vol, and the user wants the equivalent log normal (Black) vol. For consistency, we need to invert 1.2a exactly. This should be done using a global Newton method. Let us re-write 1.2a as

$$(1.5) \quad H(\sigma_B) = \sigma_N,$$

where

$$H(\sigma_B) = \sigma_B \frac{f - K}{\log f/K} \cdot \frac{1}{1 + \frac{1}{24} \left(1 - \frac{1}{120} \log^2 f/K\right) \sigma_B^2 \tau + \frac{1}{5760} \sigma_B^4 \tau^2}.$$

The problem is to find  $\sigma_B$  when the normal vol  $\sigma_N$  is given. One should start with an initial guess of

$$(1.6) \quad \sigma_B \approx \sigma_N \frac{\log f/K}{f - K} \cdot \left\{ 1 + \frac{1}{24} \left(1 - \frac{1}{120} \log^2 f/K\right) \frac{\sigma_N^2 \tau}{fK} \right\} \quad \text{if } \left| \frac{f - K}{K} \right| \geq 0.001$$

$$(1.7) \quad \sigma_B \approx \frac{\sigma_N}{\sqrt{fK}} \frac{1 + \frac{1}{24} \sigma_B^2 \tau}{1 + \frac{1}{24} \log^2 f/K} \quad \text{if } \left| \frac{f - K}{K} \right| < 0.001$$

Only one, or possibly two, Newton steps will be needed. In the Newton scheme, the derivative can be approximated by

$$H'(\sigma_B) = \frac{f - K}{\log f/K}, \quad \text{if } \left| \frac{f - K}{K} \right| \geq 0.001$$

$$H'(\sigma_B) = \frac{1}{\sqrt{fK}} \quad \text{if } \left| \frac{f - K}{K} \right| < 0.001$$

**2. Converting CEV vols to absolute or Black volatilities.** Another popular skew model is the CEV model:

$$(2.1a) \quad dR = \alpha R^\beta$$

where

$$(2.1b) \quad \beta = \text{user input CEV exponent, } 0 \leq \beta \leq 1.$$

**2.1. Converting between CEV vol and normal vol.** To convert the CEV vol  $\alpha$  into a normal (absolute) vol, one can use

$$(2.2a) \quad \sigma_N = \alpha \frac{(1 - \beta)(f - K)}{f^{1-\beta} - K^{1-\beta}} \cdot \frac{1}{1 + \frac{1 - \frac{2-2\beta+\beta^2}{120} \log^2 f/K}{1 - \frac{(1-\beta)^2}{12} \log^2 f/K} \frac{\beta(2-\beta)\alpha^2 \tau}{24(fK)^{1-\beta}}}$$

When  $f$  is very near  $K$ , or when  $\beta$  is very near 1, one needs to replace the formula with one that doesn't have the singularity at  $\beta = 1$  or  $f = K$ . To cover both possibilities, we replace the above formula with

$$(2.2b) \quad \sigma_N = \alpha (fK)^{\beta/2} \cdot \frac{1 + \frac{1}{24} \log^2 f/K}{1 + \frac{(1-\beta)^2}{24} \log^2 f/K} \cdot \frac{1}{1 + \frac{1 - \frac{2-2\beta+\beta^2}{120} \log^2 f/K}{1 - \frac{(1-\beta)^2}{12} \log^2 f/K} \frac{\beta(2-\beta)\alpha^2 \tau}{24(fK)^{1-\beta}}}$$

when

$$(2.2c) \quad (1 - \beta) \left| \frac{f - K}{K} \right| < 0.001$$

To convert the normal vol  $\sigma_N$  into a CEV vol, we should again use a global Newton method, to solve

$$(2.3a) \quad H(\alpha) = \sigma_N.$$

Here,

$$(2.3b) \quad H(\alpha) = \alpha \frac{(1 - \beta)(f - K)}{f^{1-\beta} - K^{1-\beta}} \frac{1}{1 + \frac{1 - \frac{2-2\beta+\beta^2}{120} \log^2 f/K}{1 - \frac{(1-\beta)^2}{12} \log^2 f/K} \frac{\beta(2-\beta)\alpha^2\tau}{24(fK)^{1-\beta}}}$$

if  $(1 - \beta) \left| \frac{f - K}{K} \right| \geq 0.001$

and

$$(2.3c) \quad H(\alpha) = \alpha(fK)^{\beta/2} \frac{1 + \frac{1}{24} \log^2 f/K}{1 + \frac{(1-\beta)^2}{24} \log^2 f/K} \frac{1}{1 + \frac{\beta(2-\beta)\alpha^2\tau}{24(fK)^{1-\beta}}}$$

if  $(1 - \beta) \left| \frac{f - K}{K} \right| < 0.001$

A superb initial guess is

$$(2.4) \quad \alpha \approx \frac{\sigma_N}{fK^{\beta/2}} \frac{1 + \frac{(1-\beta)^2}{24} \log^2 f/K}{1 + \frac{1}{24} \log^2 f/K} \left\{ 1 + \frac{1}{24} \frac{\beta(2-\beta)\sigma_N^2\tau}{fK} \right\}$$

As above, the derivative for Newton's method can be taken as

$$(2.5a) \quad H'(\alpha) = \frac{(1 - \beta)(f - K)}{f^{1-\beta} - K^{1-\beta}},$$

$$(2.5b) \quad H'(\alpha) = (fK)^{\beta/2}.$$

**2.2. Converting CEV vol to log normal vol.** To convert the CEV vol  $\alpha$  to a log normal (Black) vol, one should first translate it to the normal vol  $\sigma_N$ , and then use the above routine to calculate the Black vol  $\sigma_B$  from the normal vol  $\sigma_N$ .

Similarly, to convert the Black vol  $\sigma_B$  to the CEV vol  $\sigma$ , one uses the above routines to first translate it to a normal vol  $\sigma_N$ , and then translate the normal to the Black vol.



**Helpx2**

Press the <HELP>  
key twice for instant  
live assistance.

**[bloomberg.com](http://bloomberg.com)**

**Frankfurt**  
+49 69 92041 0

**Hong Kong**  
+852 2977 6000

**London**  
+44 20 7330 7500

**New York**  
+1 212 318 2000

**San Francisco**  
+1 415 912 2960

**São Paulo**  
+55 11 3048 4500

**Singapore**  
+65 6212 1000

**Sydney**  
+61 2 9777 8600

**Tokyo**  
+81 3 3201 8900

BLOOMBERG, BLOOMBERG PROFESSIONAL, BLOOMBERG MARKETS, BLOOMBERG NEWS, BLOOMBERG ANYWHERE, BLOOMBERG TRADEBOOK, BLOOMBERG BONDTRADER, BLOOMBERG TELEVISION, BLOOMBERG RADIO, BLOOMBERG PRESS and BLOOMBERG.COM are trademarks and service marks of Bloomberg Finance L.P., a Delaware limited partnership, or its subsidiaries. The BLOOMBERG PROFESSIONAL service (the "BPS") is owned and distributed locally by Bloomberg Finance L.P. (BFLP) and its subsidiaries in all jurisdictions other than Argentina, Bermuda, China, India, Japan and Korea (the "BLP Countries"). BFLP is a wholly-owned subsidiary of Bloomberg L.P. ("BLP"). BLP provides BFLP with all global marketing and operational support and service for these products and distributes the BPS either directly or through a non-BFLP subsidiary in the BLP Countries.